

**BIS-CO-SORB - bismuth subsalicylate, kaolin suspension**

**Nich Marketers, Inc**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**BIS-CO-SORB SUSPENSION**

INDICATIONS: FOR ORAL USE AS AN AID IN RELIEF OF UPSET STOMACH, NAUSEA AND DIARRHEA IN LARGE AND SMALL ANIMALS.

DO NOT USE IN CATS.

DO NOT USE IN COWS PRODUCING MILK FOR HUMAN CONSUMPTION.

**ACTIVE INGREDIENTS:**

BISMUTH SUBSALICYLATE 262 MG / 15 ML

KAOLIN 500 MG / 15 ML

IN A BASE CONTAINING: MAGNESIUM ALUMINUM SILICATE, SODIUM SACCHARIN, XANTHAN GUM, FD C RED 28, PRESERVATIVES, FLAVOR, MAGNESIUM TRISILICATE, WATER QS.

- FOR VETERINARY USE ONLY
- SHAKE WELL BEFORE USING
- KEEP OUT OF REACH OF CHILDREN
- STORE AT ROOM TEMPERATURE 69 DEGREES TO 77 DEGREES F (20 DEGREES TO 25 DEGREES C)
- PROTECT FROM FREEZING
- AVOID EXCESSIVE HEAT, OVER 104 DEGREES F (40 DEGREES C)

CAUTION: FOR VETERINARY USE ONLY

RECOMMENDED ORAL DOSAGE OF BIS-CO-SORB:

FOALS: 1 OZ (30ML) EVERY 4 HOURS. MAXIMUM 6 DOSES EVERY 24 HOURS. REPEAT AS NECESSARY, OR AS DIRECTED BY VETERINARIAN.

CALVES: 1 OZ (30ML) EVERY 4 HOURS. MAXIMUM 6 DOSES EVERY 24 HOURS. REPEAT AS NECESSARY, OR AS DIRECTED BY VETERINARIAN.

SHEEP AND GOATS: 1/2 OZ (15ML) EVERY 4 HOURS. MAXIMUM 6 DOSES EVERY 24 HOURS. REPEAT AS NECESSARY, OR AS DIRECTED BY VETERINARIAN.

DOGS: 1 TEASPOON (5ML) EVERY 4 HOURS. MAXIMUM 6 DOSES EVERY 24 HOURS. REPEAT AS NECESSARY, OR AS DIRECTED BY VETERINARIAN.

HOGS: 2 OZ (60ML) PER 100 LBS OF BODY WEIGHT EVERY 3 TO 4 HOURS. MAXIMUM 6 DOSES EVERY 24 HOURS. REPEAT AS NECESSARY, OR AS DIRECTED BY VETERINARIAN.

NOTE: DOSAGE SHOULD BE ADJUSTED AS NECESSARY.

ALLOW ANIMAL TO HAVE ACCESS TO CLEAN FRESH DRINKING WATER AS NEEDED.

WARNING: IF CLINICAL SIGNS PERSIST AFTER 2 DAYS, RECONSULT A VETERINARIAN.

WARNING: CONTAINS SALICYLATES

**BIS-CO-SORB Suspension 1 gallon (42065-222-01)**

NDC 42065-222-01

BIS-CO-SORB  
SUSPENSION

BISMUTH SUBSALICYLATE AND KAOLIN SUSPENSION

FOR VETERINARY USE ONLY

ORAL MEDICINE FOR USE AS AN AID IN RELIEF OF UPSET STOMACH, AS AN  
ANTIDIARRHEAL AID, AND TO AID IN RELIEF OF NAUSEA

FOR ORAL USE IN CALVES, FOALS, SHEEP, GOATS, HOGS, AND DOGS

NET CONTENTS:  
1 GALLON (128 FL OZ)

MANUFACTURED FOR:  
NICH MARKETERS, INC.  
GULF BREEZE FL 32563

**INDICATIONS:** FOR ORAL USE AS AN AID IN  
RELIEF OF UPSET STOMACH, NAUSEA, AND DIAR-  
RHEA IN LARGE AND SMALL ANIMALS.  
DO NOT USE IN CATS.  
DO NOT USE IN COWS PRODUCING MILK FOR  
HUMAN CONSUMPTION.

**ACTIVE INGREDIENTS:**  
BISMUTH SUBSALICYLATE 262 mg / 15 ml  
KAOLIN 500 mg / 15ml  
IN A BASE CONTAINING: MAGNESIUM ALUMINUM  
SILICATE, SODIUM SACCHARIN, XANTHAN GUM,  
FD&C RED #28, PRESERVATIVES, FLAVOR, MAGNE-  
SIUM TRISILICATE, WATER QS.

**FOR VETERINARY USE ONLY**  
**KEEP OUT OF REACH OF CHILDREN**  
**SHAKE WELL BEFORE USING**  
**STORE AT ROOM TEMPERATURE**  
**69° TO 77°F (20° TO 25°C)**  
**PROTECT FROM FREEZING**  
**AVOID EXCESSIVE HEAT**  
**OVER 104°F (40°C)**

NDC#42065-222-01

**BIS-CO-SORB**  
**S U S P E N S I O N**

**BISMUTH SUBSALICYLATE AND  
KAOLIN SUSPENSION**

**FOR VETERINARY USE ONLY**

ORAL MEDICINE FOR USE AS AN AID IN  
RELIEF OF UPSET STOMACH, AS AN  
ANTIDIARRHEAL AID, AND TO AID IN RELIEF  
OF NAUSEA  
FOR ORAL USE IN CALVES, FOALS, SHEEP,  
GOATS, HOGS, AND DOGS

**Net contents:**  
**1 Gallon (128 fl. oz.)**



MANUFACTURED FOR:  
NICH Marketers, Inc.  
Gulf Breeze, FL 32563

**CAUTION: FOR VETERINARY USE ONLY**

**RECOMMENDED ORAL DOSAGE OF BIS-CO-SORB:**  
Foals: 1 oz (30 ml) every 4 hours. Maximum 6 doses every 24  
hours. Repeat as necessary, or as directed by veterinarian.  
Calves: 1 oz (30 ml) every 4 hours. Maximum 6 doses every  
24 hours. Repeat as necessary, or as directed by veterinarian.  
Sheep & Goats: 1/2 oz (15 ml) every 4 hours. Maximum 6  
doses every 24 hours. Repeat as necessary, or as directed by  
veterinarian.  
Dogs: 1 teaspoon (5 ml) every 4 hours. Maximum 6 doses  
every 24 hours. Repeat as necessary, or as directed by veteri-  
narian.  
Hogs: 2 oz (60 ml) per 100 lbs of body weight every 3 to 4  
hours. Maximum 6 doses every 24 hours. Repeat as neces-  
sary, or as directed by veterinarian.

**NOTE:** DOSAGE SHOULD BE ADJUSTED AS NECESSARY.  
ALLOW ANIMAL TO HAVE ACCESS TO CLEAN FRESH DRINKING  
WATER AS NEEDED.

**WARNING:** If clinical signs persist after 2 days, reconsult a  
veterinarian.

**WARNING:** CONTAINS SALICYLATES.  
DO NOT USE IN CATS.  
DO NOT USE IN COWS PRODUCING MILK FOR HUMAN CONSUMPTION.

5-2008

**BIS-CO-SORB Suspension 60ml (42065-222-02)**

NDC 42065-222-02

BIS-CO-SORB  
SUSPENSION

BISMUTH SUBSALICYLATE AND KAOLIN SUSPENSION

FOR VETERINARY USE ONLY

60 ml.

EASY

DOSE

SYRINGE

MANUFACTURED FOR:  
NICH MARKETERS, INC.  
GULF BREEZE FL 32563

NDC# 42065-222-02

# BIS-CO-SORB

## SUSPENSION

60 ml.  
EASY  
DOSE  
SYRINGE

**BISMUTH SUBSALICYLATE AND KAOLIN SUSPENSION  
FOR VETERINARY USE ONLY.**

**INDICATIONS:** FOR ORAL USE AS AN AID IN RELIEF OF UPSET STOMACH, NAUSEA, AND DIARRHEA IN LARGE AND SMALL ANIMALS.

**ACTIVE INGREDIENTS:**

BISMUTH SUBSALICYLATE 262 mg / 15 ml

KAOLIN 500 mg / 15ml

IN A BASE CONTAINING: MAGNESIUM ALUMINUM SILICATE, SODIUM SACCHARIN, XANTHAN GUM, FD&C RED #28, PRESERVATIVES, FLAVOR, MAGNESIUM TRISILICATE, WATER QS.

**RECOMMENDED ORAL DOSAGE OF BIS-CO-SORB:**

**Sheep & Goats:** 1/2 oz (15 ml) every 4 hours. Maximum 6 doses every 24 hours. Repeat as necessary, or as directed by veterinarian.

**Dogs:** 1 teaspoon (5 ml) every 4 hours. Maximum 6 doses every 24 hours. Repeat As necessary, or as directed by veterinarian.

**NOTE:** DOSAGE SHOULD BE ADJUSTED AS NECESSARY. ALLOW ANIMAL TO HAVE ACCESS TO CLEAN FRESH DRINKING WATER AS NEEDED.

**WARNING:** If clinical signs persist after 2 days, reconsult a veterinarian.

**WARNING:** CONTAINS SALICYLATES. DO NOT USE IN CATS. DO NOT USE IN COWS PRODUCING MILK FOR HUMAN CONSUMPTION.

**KEEP OUT OF REACH OF CHILDREN  
SHAKE WELL BEFORE USING**

**STORE AT ROOM TEMPERATURE 69° TO 77°F (20° TO 25°C)  
PROTECT FROM FREEZING AVOID EXCESSIVE HEAT OVER 104°F (40°C)**



**MANUFACTURED FOR:  
NICH MARKETERS, INC.  
GULF BREEZE, FL 32563**

5-2008

## BIS-CO-SORB

bismuth subsalicylate, kaolin suspension

### Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:42065-222
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1H8YE)		BISMUTH SUBSALICYLATE	262 mg in 15 mL	
KAOLIN (UNII: 24H4NWX5CO) (KAOLIN - UNII:24H4NWX5CO)		KAOLIN	500 mg in 15 mL	
Inactive Ingredients				
Ingredient Name			Strength	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)				
SACCHARIN SODIUM DIHYDRATE (UNII: SB8ZUX40TY)				
XANTHAN GUM (UNII: TTV12P4NEE)				
D&C RED NO. 28 (UNII: 767IP0Y5NH)				
MAGNESIUM TRISILICATE (UNII: C2E1CI501T)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42065-222-01	3785 mL in 1 BOTTLE		
2	NDC:42065-222-02	60 mL in 1 SYRINGE		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other			02/23/2012	

**Labeler** - Nich Marketers, Inc (883363012)

**Registrant** - Performance Products, Inc. (006265888)

## Establishment

Name	Address	ID/FEI	Business Operations
Performance Products, Inc.		006265888	manufacture

## Establishment

Name	Address	ID/FEI	Business Operations
5N Plus Lübeck GmbH		316594993	api manufacture